

WHAT IS CLAIMED IS:

sub
or
consisting of:

An isolated nucleic acid comprising a member selected from the group

- (a) a polynucleotide having at least 75% sequence identity, as determined by the GAP algorithm under default parameters, to a polynucleotide of SEQ ID NO: 1;
- (b) a polynucleotide encoding a polypeptide of SEQ ID NO: 2;
- (c) a polynucleotide amplified from a nucleic acid library using primers which selectively hybridize, under stringent hybridization conditions, to loci within a polynucleotide of SEQ ID NO: 1;
- (d) a polynucleotide which selectively hybridizes, under stringent hybridization conditions and a wash in 0.1X SSC at about 60 to 65°C, to a polynucleotide of SEQ ID NO: 1;
- (e) a polynucleotide of SEQ ID NO: 1;
- (f) a polynucleotide which is complementary to a polynucleotide of (a), (b), (c), or (e); and
- (g) a polynucleotide comprising at least 50 contiguous nucleotides from a polynucleotide of (a), (b), (c), (d), (e), or (f).

2. A recombinant expression cassette, comprising a member of claim 1 operably linked, in sense or anti-sense orientation, to a promoter.

3. A host cell comprising the recombinant expression cassette of claim 2.

sub
or
4. A transgenic plant comprising a recombinant expression cassette of claim 2.

5. The transgenic plant of claim 4, wherein said plant is a monocot.

6. The transgenic plant of claim 4, wherein said plant is selected from the group consisting of: maize, soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton, rice, barley, and millet.

5 7. A transgenic seed from the transgenic plant of claim 4.

Sub
as 8. A method of modulating the level of nitrate-responsive root transcriptional factor in a plant, comprising:

- 10 (a) introducing into a plant cell a recombinant expression cassette comprising a root transcriptional factor polynucleotide of claim 1 operably linked to a promoter;
- (b) culturing the plant cell under plant cell growing conditions;
- (c) regenerating a plant from said plant cell; and
- 15 (d) inducing expression of said polynucleotide for a time sufficient to modulate the level of nitrate-responsive root transcriptional factor in said plant.

9. The method of claim 8, wherein the plant is maize.

20 10. An isolated protein comprising a member selected from the group consisting of:

- (a) a polypeptide of at least 20 contiguous amino acids from a polypeptide of SEQ ID NO: 2;
- (b) a polypeptide of SEQ ID NO: 2;
- 25 (c) a polypeptide having at least 75% sequence identity to, and having at least one epitope in common with, a polypeptide of SEQ ID NO: 2, wherein said sequence identity is determined by the GAP algorithm under default parameters; and,
- (d) at least one polypeptide encoded by a member of claim 1.